

Ab Initio Treatment of Electron Correlations in Polymers: Lithium Hydride Chain and Beryllium Hydride Polymer

A. Abdurahman, A. Shukla, M. Dolg

published in

Modern Methods and Algorithms of Quantum Chemistry,
J. Grotendorst (Ed.), John von Neumann Institute for Computing,
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 28, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

Ab initio treatment of electron correlations in polymers: lithium hydride chain and beryllium hydride polymer

Ayjamal Abdurahman, Alok Shukla and Michael Dolg

*Max-Planck-Institut für Physik komplexer Systeme,
Nöthnitzer Str. 38, D-01187 Dresden, Germany*

Correlated *ab initio* electronic structure calculations are reported for the polymers lithium hydride chain $[LiH]_{\infty}$ and beryllium hydride $[Be_2H_4]_{\infty}$. First, employing a Wannier-function-based approach, the systems are studied at the Hartree-Fock level, by considering chains, simulating the infinite polymers. Subsequently, for the model system $[LiH]_{\infty}$, the correlation effects are computed by considering virtual excitations from the occupied Hartree-Fock Wannier functions of the infinite chain into the complementary space of localized unoccupied orbitals, employing a full-configuration-interaction scheme. For $[Be_2H_4]_{\infty}$, however, the electron correlation contributions to its ground state energy are calculated by considering finite clusters of increasing size modelling the system. Methods such as Møller–Plesset second-order perturbation theory and coupled-cluster singles, doubles and triples level of theory were employed. Equilibrium geometry, cohesive energy and polymerization energy are presented for both polymers, and the rapid convergence of electron correlation effects, when based upon a localized orbital scheme, is demonstrated.